



**Udbytte i hkg/ha korrigeret til 85 % tørstof**

NS Abildgård		Varde		Brande		Tystofte		Koldkærgård		999		rkflge
113	KWS Erebor	104	KWS-H273	102	KWS-H273	112	KWS-H273	108	KWS Erebor	105	KWS-H273	1
132	Gns af sorter	111.4	Blanding	119.1	Blanding	106	KWS-H274	108	Gns af sorter	103	Gns af sorter	2
108	KWS-H258	99	KWS Falkor	99	KWS-H274	105	KWS Erebor	103	KWS-H273	103	KWS Erebor	3
108	KWS Falkor	99	KWS-H274	98	KWS-H258	127	Gns af sorter	102	KWS-H258	102	KWS-H274	4
108	KWS-H274	99	KWS-H258	98	KWS Falkor	102	KWS-H258	84.6	Blanding	102	KWS-H258	5
107	KWS-H273	96	KWS Erebor	95	KWS Erebor	102.9	Blanding	99	KWS-H274	103.4	Blanding	6
99.1	Blanding	127	Gns af sorter	134	Gns af sorter	89	KWS Falkor	92	KWS Falkor	97	KWS Falkor	7
9	LSD 0.05	11	LSD 0.05	10	LSD 0.05	8	LSD 0.05	6	LSD 0.05	4	LSD 0.05	

## Translations

Afgrødehøjde	<i>Crop height</i>	Standardkvalitet	<i>Standard quality</i>
Blomstring	<i>Flowering</i>	Stivelsesindhold	<i>Starch content</i>
Blødgøring	<i>Softening</i>	Strå længde, cm	<i>Straw length</i>
Brunrust	<i>Brown rust (Puccinia recondita)</i>	Udbytte	<i>Yield</i>
Brødhøjde	<i>Bread height</i>	Vandoptagelse	<i>Water absorption</i>
Brødvolumen	<i>Bread volume</i>		
Bygrust	<i>Barley Rust (Puccinia hordei)</i>		
Dyrkningsegenskaber	<i>Agronomic traits</i>		
EFOSi	<i>Enzyme digestible organic matter at ileum</i>		
EFOSsvin	<i>Enzyme digestible organic matter in pigs</i>		
Erucasyre	<i>Erucic acid</i>		
FEso pr. hkg	<i>Feed units, adult pigs</i>		
FEsv pr. hkg	<i>Feed units, growing pigs</i>		
fht	<i>Index</i>		
Faldtal	<i>Falling number</i>		
Foderkvalitet	<i>Feed quality</i>		
Frøkvalitet	<i>Seed quality</i>		
Frøvægt	<i>Seed weight</i>		
Glucosinolatindhold	<i>Glucosinolate content</i>		
Gluten i kerner (14% vand)	<i>Gluten content in grains at 14 % water</i>		
Gns.	<i>Average</i>		
Gråplet/brunplet	<i>Septoria tritici/Stagonospora nodorum</i>		
Gulrust	<i>Yellow rust (Puccinia striiformis)</i>		
hkg/ha korrigeret til 85 % tørstof	<i>hkg/ha adjusted to 85% dry matter</i>		
Hvedebladplet	<i>Tan spot, DTR (Pyrenophora tritici-repentis)</i>		
Karakter	<i>Score</i>		
Kernekvalitet	<i>Grain quality</i>		
Klæbrighed	<i>Stickyness</i>		
Kornvægt, mg pr. korn	<i>Thousand kernel weight (mg/kg)</i>		
Kvalitetsegenskaber	<i>Quality traits</i>		
Led	<i>Entry</i>		
Lejesæd	<i>Lodging</i>		
Linolénysyre	<i>Linolenic acid</i>		
Linolsyre	<i>Linoleic acid</i>		
Meldug	<i>Mildew (Erysiphe graminis)</i>		
Meludbytte	<i>Flour yield</i>		
Modning, dato	<i>Ripeningdate</i>		
Nedknækning, aks	<i>Necking</i>		
Nedknækning, strå	<i>Brackling</i>		
Observations-parceller	<i>Observation-plots</i>		
Olieindhold	<i>Oil content</i>		
Oliesyre	<i>Oleic acid</i>		
Plantehøjde	<i>Plant height</i>		
Proteinindhold, pct.	<i>Protein content</i>		
Ramularia	<i>Ramularia (Ramularia collo-cygni)</i>		
Rumvægt, g pr. liter	<i>Specific weight</i>		
Sedimentation	<i>Zeleny sedimentation value</i>		
Skala	<i>Scale</i>		
Skoldplet	<i>Leaf Blotch (Rhynchosporium secalis)</i>		
Sort	<i>Variety</i>		
Sort., pct. kerner > 2,5 mm	<i>Grading, pct. kernels &gt; 2.5 mm</i>		
Sort., pct. kerner > 2,8 mm	<i>Grading, pct. kernels &gt; 2.8 mm</i>		
Sortskode	<i>Variety code</i>		
Stabilitet	<i>Stability</i>		